



Forage selection by Masai giraffes (*Giraffa camelopardalis tippelskirchi*) at multiple spatial scales

MATANA LEVI,^{1,*} DEREK E. LEE,² MONICA L. BOND,³ AND ANNA C. TREYDTE^{1,4,5}

¹Department of Sustainable Agriculture, Biodiversity and Ecosystem Management, School of Life Sciences and Bioengineering (LiSBE), Nelson Mandela African Institution of Science and Technology, P.O. Box 447, Arusha, Tanzania

²Department of Biology, Pennsylvania State University, University Park, PA 16802, USA

³Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich 8057, Switzerland

⁴Ecology of Tropical Agricultural Systems, Hans-Ruthenberg Institute, University of Hohenheim, Stuttgart 70599, Germany

⁵Department of Physical Geography, Stockholm University, Stockholm 10691, Sweden

*To whom correspondence should be addressed: levi.matana@gmail.com

Management of rangelands requires knowledge of forage species that are preferred or avoided by wildlife and livestock. A recent expansion of woody vegetation into previously open grasslands in African savanna ecosystems negatively impacts many mammalian grazers. Nevertheless, the ecological role of bush encroacher plant species as food may present a benefit for browsing species. We quantified diet selection by Masai giraffes (*Giraffa camelopardalis tippelskirchi*) through foraging observations and vegetation sampling in the Tarangire Ecosystem of Tanzania, which includes large areas of a native shrub that livestock managers have classified as an encroacher species (*Dichrostachys cinerea*). We compared woody plant species used by giraffes for foraging with availability at two different spatial scales during the wet and dry seasons. Giraffes selected some woody plants such as *Vachellia* species while significantly avoiding others, both at the local and landscape scales. Giraffes preferred foraging on *D. cinerea* at both spatial scales and in both the wet and dry seasons. Management that has focused on benefiting grazing livestock by removal of encroaching species (e.g., *D. cinerea*) may have unintended consequences for wildlife, especially for browsing species like giraffes that feed extensively on the expanding bush species.

Key words: browser, rangeland ecology, resource selection, savanna landscapes, Tanzania, woody vegetation

Understanding resource selection by animals has been a cornerstone of basic ecology and rangeland management for decades (Pellew 1983; Mahenya et al. 2016). Resource selection studies provide empirical evidence about feeding ecology and habitat suitability, which can inform species conservation (Pellew 1984a, 1984b; Arthur et al. 1996; Manly et al. 2002; Bryson-Morrison et al. 2017). Recent and rapid transformation of habitat by humans has led to increased concerns about proper management of rangelands (Belayneh and Tessema 2017; Birhane et al. 2017; Devine et al. 2017). In African savannas, this transformation is sometimes expressed as an expansion of woody vegetation into open grass-dominated lands, which holds the potential to alter ecosystem processes (Hudak and Wessman 1998; Van de Vijver et al. 1999; Ludwig 2001; Roques et al. 2001; Ludwig et al. 2008; Devine et al. 2017). Savannas support the highest densities of wild mammalian

herbivores of any biome (Grady and Hoffmann 2012; Shorrock and Bates 2015), so understanding how herbivores select food resources is critical for maintaining ecosystem function and for conserving endangered species such as the Masai giraffe (*Giraffa camelopardalis tippelskirchi*).

The Masai giraffe is an endangered megaherbivore inhabiting savanna ecosystems of East Africa (Bolger et al. 2019). Abundance of Masai giraffes has declined 49–51% during the last three decades (Bolger et al. 2019), primarily because of habitat loss and degradation associated with human activities such as deforestation, livestock grazing, bush fires, and illegal hunting (Zarovali et al. 2007; Belayneh and Tessema 2017; Bolger et al. 2019). Forage species most commonly used by giraffes in East African savannas include *Vachellia* spp., *Commiphora* spp., and *Combretum* spp. (Pellew 1983, 1984b; Mahenya et al. 2016); however, encroaching woody

plants may serve as alternative forage for browsing giraffes. *Dichrostachys cinerea* is a native shrub that can form clonal mats in areas heavily impacted by domestic livestock grazing (Tjelele et al. 2014), and some land managers believe that a reduction of *D. cinerea* is needed to maintain grazing resources for wildlife and livestock (Njagi 2019). Studies have examined the response of grazing mammalian species to this shift from grass-dominated to woody-dominated vegetation dynamics (Dalle et al. 2006; Smit and Prins 2015), but little is known about how this shift might affect browsing species such as giraffes. We aimed to quantify forage selection by Masai giraffes in a human-influenced landscape with areas dominated by the encroaching woody species, *D. cinerea*.

Resource selection is defined as the ratio of proportional use over availability for a given plant species or taxa (Johnson 1980; Manly et al. 2002; Dumont et al. 2007). In theory, animals should select forage of the highest quality in order to meet their nutritional requirements (Pellew 1984a; Shipley 2007). Selection of forage resources occurs at different spatial scales, ranging from an entire geographic area (landscape) exploited by a species to the selection of forage within foraging patches (Johnson 1980; Johnson et al. 2002; Boyce 2006). Different spatial scales affect the proportions of available forage resources and, consequently, resource selection (Bissonetie et al. 1997). Thus, inferences made during resource selection studies and associated management decisions must account for the spatial scale of selection being considered (Johnson 1980; Wiens 1981; Orians and Wittenberger 1991; Manly et al. 2002). Using only one spatial scale of resource assessment, especially in heterogeneous landscapes, is potentially biased (Arthur et al. 1996) and may misinform rangeland management decisions (Wiens 1981; Kotliar and Wiens 2013).

We examined diet and forage selection at local and landscape scales by Masai giraffes in the Tarangire Ecosystem of northern Tanzania. We gave particular attention to the use of *D. cinerea* by giraffes because we were interested in determining if this bush species, which often is classified as an encroaching species, is an important forage species for giraffes. We posed six research questions: (i) What woody plant species do giraffes eat in the Tarangire Ecosystem? (ii) Is the use of forage species random or selective? (iii) If selective, which species are preferred or avoided? (iv) To what extent is *D. cinerea* an important component in giraffe diets? (v) Does forage selection of giraffes differ between the wet and the dry season? (vi) Does forage selection of giraffes differ between landscape and foraging patch spatial scales?

Giraffes live in a fission/fusion system, with herd sizes averaging four to six individuals but ranging upwards of 60 individuals (Bond et al. 2019). Giraffe herds can be single sex or mixed groups comprised of individuals feeding on an array of forage plants (Bercovitch and Berry 2013; Bond et al. 2019). We predicted that giraffes would be selective in their foraging, as demonstrated in previous studies in other ecosystems (e.g., Transvaal, South Africa: Sauer et al. 1977; Sahel, Niger: Caister et al. 2003; South Africa: Parker et al. 2003) and that foraging selection would change according to spatial scale (Bissonetie et al. 1997) and season (Sauer et al. 1977; Berry and Bercovitch

2016). We also predicted that giraffes would avoid browsing on *D. cinerea*, a shrub that is believed to be unpalatable to large mammalian herbivores. Lastly, we predicted that local-scale forage selection would be less visible than the selection at the landscape scale because, while foraging, giraffes would have already chosen to be in locations with their preferred food sources (Bissonetie et al. 1997; Anderson et al. 2005).

MATERIALS AND METHODS

Study area.—We conducted our study in the Tarangire Ecosystem, northern Tanzania (Fig. 1; between 35°80′, –3°80′ to 36°20′, –4.20′). The Tarangire Ecosystem receives a mean rainfall of 529 mm (Peterson 1978), distributed in two periods called short rains (October–January) and long rains (February–May) separated by a dry (June–September) season (Galanti et al. 2006). Its undulating plateaus are composed mainly of dark-red sandy clay loam, waterlogged areas, and floodplains of black cotton soils, with elevation varying between 900 and 1,200 m above sea level (Kahurananga and Silkiluwasha 1997; Galanti et al. 2006). Our study area in the Tarangire Ecosystem included a national park (Tarangire National Park) and rangelands outside the park (Manyara Ranch and Randilen Wildlife Management Area), where suitable giraffe habitat exists. The study area was predominantly savanna, comprised of open grasslands, woodlands, riverine forests, and shrublands, and falls in the semiarid zone, based on rangeland classification by Pratt et al. (1966). The study area represents one of the largest remaining refugia during the dry season for migratory ungulates in the country (Stoner et al. 2007; Bolger et al. 2008) as well as an important giraffe metapopulation (Lee and Bolger 2017). During the wet season, as food resources become abundant, wild herbivores were distributed widely across this human-influenced landscape (Newmark et al. 1994; Galanti et al. 2006; Msoffe et al. 2011).

Behavioral observations.—Foraging observations were collected during 15 days per month in March and April 2019 (wet season observations), and August and September 2019 (dry season observations). Each day, we drove in a car along the road network of our study area during daylight hours at a speed of 5–20 km/h, searching for giraffes. At an opportunistic encounter of a giraffe herd, we started 2-h observation records. During each 2-h observation period, we obtained foraging observations using instantaneous scan sampling (Martin and Bateson 1993). We conducted 5-min scans at 10-min intervals, using binoculars (10 × 50), with one observer and one data recorder. Foraging events were recorded from the left to the right-hand side of the group in a first seen–first recorded style within 5 min (Martin and Bateson 1993). In every scan, we identified and recorded each plant species eaten by each foraging giraffe within the group to assess general diet composition. Giraffes were followed as closely as possible without disturbing them, at a distance of 60–200 m. We considered a herd as the sampling unit, so we summarized individual foraging data during an observation period into proportional use by the entire herd (Fig. 1). The problem of group fusion was avoided by maintaining the original number of individuals that




Fig. 1.—Map showing systematic vegetation point transects (black and white stars), where landscape-scale vegetation sampling was conducted, and the locations of Masai giraffe foraging observations, where local-scale vegetation sampling was conducted during the dry (gray squares) and the wet season (black filled circles) in the Tarangire Ecosystem, in 2019.

our scans had started with. In the case of group fission, we tried as much as possible to continue observing the same individuals and treating scattered individuals as a group. We ended observations when some members of the group were no longer within our vicinity. Observations of giraffes followed ASM guidelines (Sikes et al. 2016), and all work was permitted by the Tanzania Commission for Science and Technology (COSTECH) and the Tanzania Wildlife Research Institute (TAWIRI).

Forage availability.—We assessed availability of forage plant species at local (within foraging patches) and landscape (entire study area) scales. Woody plants were identified directly in the field by a botanist while unidentified woody plant specimens were pressed in a plant press, assigned a collection number (symbol), and recorded in a field notebook for further identification. At both scales, our assessment was aimed at capturing proportions of available species considering both used and nonused forage resources (Bissonetie et al. 1997). We collected data on available woody plants in March and April 2019 (wet season observations) and in August and September 2019 (dry season observations).

The local scale was intended to assess forage availability at a fine-scale of selection within areas where giraffes were observed foraging (Johnson 1980; Manly et al. 2002; Boyce 2006). To quantify the availability of woody plant species at local scale, we sampled vegetation plots along giraffe routes at patches where giraffes had been observed foraging, immediately after the foraging herd had moved on. In each foraging patch, we laid three circular plots of 10 m in diameter, 5 m away from each other. Within each circular plot (Banda et al. 2008; Chytry et al. 2013), we identified all woody species and determined the percent cover for each forage species.

To quantify forage availability at the landscape scale, we conducted systematic vegetation sampling within the area where giraffes had been seen regularly for the last 6 years (Lee and Bond 2016; Lee et al. 2016; Lee and Bolger 2017). We established 44 points systematically across the landscape using QGIS 2.18.12; all points were 5 km apart. At each point, we established four “strip transects” of 40 × 10 m to the North, East, South, and West (Lindgren and Sullivan 2001). Within each strip transect, we recorded the woody vegetation following the same three-circular-plot methodology as used for the local vegetation assessment.

Data analyses.—To understand which plants were selected by giraffes, we listed all woody plant species giraffes were observed to consume. Our selection analysis applied a chi-square test in combination with Bonferroni confidence intervals (CIs; Neu et al. 1974) for both local and landscape scales. To determine whether foraging was selective or random and if giraffe forage selection differed with regard to spatial scale and season, we included in our analysis those forage species with >5 forage observations (Manly et al. 2002). We used a chi-square test to compare observed foraging observations against the expected foraging observations (calculated based on proportions of forage availability).

To determine patterns of selection of forage at both spatial scales and across the seasons, we used Manly’s selection ratio (\hat{W}) and Bonferroni CIs (Neu et al. 1974; Manly et al. 2002). Manly’s selection ratios for each forage species were calculated based on proportions of the forage resource used versus its availability (Manly et al. 2002). To obtain proportions of forage used, P_u , for each foraged woody plant species, we

calculated the number of foraging observations in each consumed species divided by the total number of foraging observations for all consumed species across all giraffe herds. We computed local-scale proportional availability (local P_a) of all woody plant species for the foraging route taken by each herd using each species' average percent cover across the three circular plots. We obtained landscape-scale available proportions (landscape P_a) for each woody plant species by taking the average proportion of occurrence of all woody plant species from the systematic vegetation sampling. We determined preference or avoidance as \hat{W} being greater or less than 1, respectively. Significance of selection for a given species was based on the CI of each \hat{W} : if the CI was >1 , this was categorized as preferred, and if the CI was <1 the species was categorized as avoided, while no selection occurred if the CI of \hat{W} included 1 (Manly et al. 2002). We used the generic name, *Vachellia*, in place of the former genus *Acacia* for Africa and Asia as agreed by the Nomenclature Session of the Seventeenth International Botanical Congress (IBC) in 2003 (Maslin 2008).

RESULTS

Diet composition and forage selection

The instantaneous scan sampling produced 3,728 foraging observations during 1,250 scans of 105 giraffe herds. In our vegetation sampling at the landscape scale ($n = 44$) and the local scale ($n = 105$), we identified 118 woody plant species. Giraffes consumed 38 of the 118 woody plant species that we identified, with 29 consumed in the wet and 33 in the dry season (Supplementary Data SD1 and SD2). The most-consumed plant species in both the wet and dry season were *Vachellia tortilis* (25%), *D. cinerea* (23%), *V. mellifera* (17%), *V. drepanolobium* (9%), *Balanites aegyptiaca* (7%), *V. kirkii* (4%), *Dalbergia melanoxylon* (4%), *Maerua triphylla* (2%), and *Ziziphus mucronata* (1%).

Giraffes foraged nonrandomly at both the local ($\chi^2 = 1,068$, d.f. = 30, $P < 0.001$) and landscape scales ($\chi^2 = 1,670$, d.f. = 29, $P < 0.001$). Giraffes preferred *V. tortilis*, *V. mellifera*, and *V. drepanolobium* consistently through the year (Fig. 2; Supplementary Data SD3 and SD4). In contrast to what we had predicted, giraffes exhibited a high and year-round preference for the native shrub *D. cinerea* at both local and landscape scales (Fig. 2; Supplementary Data SD3, SD4 and SD5). Giraffes also avoided some woody species such as *Kigelia africana*, *Commiphora schimperi*, *Commiphora africana*, *Adansonia digitata*, and *Euphorbia candelabrum* (Fig. 2; see Supplementary Data SD3 and SD4).

Local scale.—Giraffes were selective in foraging at the local spatial scale ($\chi^2 = 1,068$, d.f. = 30, $P < 0.001$), which was visible in both dry ($\chi^2 = 709$, d.f. = 28, $P < 0.001$) and wet seasons ($\chi^2 = 501$, d.f. = 21, $P = 0.001$). Giraffes demonstrated a strong preference toward *D. cinerea*, *V. tortilis*, *V. drepanolobium*, and *V. kirkii* at the local scale during the wet season, whereas during the dry season, *D. cinerea*, *V. mellifera*, *Strychnos potatorum*, and *Combretum zeyheri* were selected more frequently than their proportional availability (Fig. 2; Supplementary Data




Fig. 2.—Local- (A and B) and landscape-scale (C and D) forage selection ratios and direction of forage selection indicated by confidence intervals ($\hat{W} \pm CIs$) for the 15 most frequently selected forage species by Masai giraffes during both the wet (A and C) and dry season (B and D) in the Tarangire Ecosystem, in 2019. Shaded bars = significantly preferred, black bars = significantly avoided, and open bars = not preferred/avoided. The black dotted line (x-axis) indicates the threshold for forage selectivity. The selection coefficient is significant if the CI for \hat{W} does not contain the value 1.

SD3). Giraffes used *B. aegyptiaca*, *Carrisa spinorum*, *Scolopia zeyheri*, and *Z. mucronata* in proportion with their availability, but avoided *A. digitata*, *Capparis fascicularis*, *Cordia monoica*, *C. schimperii*, *E. candelabrum*, *K. africana*, *Lanchoarpus eriocalyx*, *V. nilotica*, and *V. senegal* in both the wet and dry seasons at the local scale (Fig. 2; Supplementary Data SD3).

Landscape scale.—Giraffes also foraged selectively at the landscape scale ($\chi^2 = 1,670$, d.f. = 29, $P < 0.001$), during both the dry ($\chi^2 = 1,495$, d.f. = 29, $P < 0.0001$) and wet ($\chi^2 = 856$, d.f. = 29, $P < 0.0001$) seasons. At the landscape scale, *B. aegyptiaca*, *D. cinerea*, *V. drepanolobium*, *V. mellifera*, and *V. tortilis* were the most strongly favored forage species throughout the year. During the wet season, *B. aegyptiaca*, *D. cinerea*, *V. drepanolobium*, *V. mellifera*, *V. tortilis*, and *V. kirkii* were preferred, whereas in the dry season, giraffe preferred foraging on *B. aegyptiaca*, *D. cinerea*, *Ficus natalensis*, *M. triphylla*, *V. drepanolobium*, *V. mellifera*, and *V. tortilis* relative to availability on the landscape (see Fig. 2 and Supplementary Data SD4).

DISCUSSION

The diet of Masai giraffes in the Tarangire Ecosystem consisted of a variety of woody plant species and confirmed previous studies in other regions (Pellew 1984b; Parker and Bernard 2005; Dagg 2014; Berry and Bercovitch 2016) that giraffes will forage on many plant species but concentrate on a narrow range of forage options, most notably *Vachellia* species. For example, in the Serengeti National Park, five forage species accounted for about 70% of the giraffe's diet (Pellew 1984b). Parker and Bernard (2005) assessed the giraffe diet in the Eastern Cape Province of South Africa, where 46 forage species were consumed, but *Vachellia karroo* and *Rhus longispina* comprised 60% of the total diet. In a recent study by Mahenya et al. (2016) in Arusha National Park in Tanzania, the giraffes' diet was 90% composed of *V. xanthophloea*. Similarly, in our study in Tarangire Ecosystem, *D. cinerea* and *Vachellia* species contributed about 65% of the overall giraffe diet. Selection for *D. cinerea* and *Vachellia* species was likely due to the high nutritional value and digestibility of these woody plant species (Sauer 1983; Pellew 1984a). Giraffes also demonstrated strong avoidance of some woody species such as *K. africana* and *A. digitata*, likely because of their poor digestibility (Woodward and Coppock 1995; Proll et al. 2018).

We detected seasonal differences in forage selection by Masai giraffes for particular plant species at both the local and landscape scale, probably because of a decrease in forage plant availability during the dry season compared to the wet season (Beyer and Hauffer 1994; Whittingham et al. 2005; Boyce 2006). Masai giraffes appear to demonstrate seasonal forage use toward a few species to guarantee sufficient amount and nutrient content of food (Sauer et al. 1977; Sauer 1983; Pellew 1984b; Parker et al. 2003). Most deciduous woody plants such as *C. zeyheri*, *D. cinerea*, *V. kirkii*, *V. mellifera*, and *V. tortilis* contributed strongly to giraffe diet during the wet season, but some became less important in the dry season, in agreement with similar studies (Sauer 1983; Pellew 1984b).

Giraffe foraging preferences in the Tarangire Ecosystem switched to semideciduous plants during the dry season, most notably *Combretum* species, which retain their leaves and protein content as the dry season progresses (Sauer 1983). Our results are in line with those of Sauer (1983) that the selection of *Vachellia* species often declines in the dry season because of a decrease in protein and water content, while *Combretum* species lose their proteins more slowly, making them a suitable forage source for longer into the dry season. This result likely explains why giraffe groups in the Tarangire Ecosystem were larger in stands of *Combretum* during the dry compared with the long rainy season (Bond et al. 2019).

Our results did not support our prediction that local-scale forage selectivity would be expressed less strongly than at the landscape scale. However, giraffe forage preferences were sensitive to spatial scale changes (Bissonetie et al. 1997; Anderson et al. 2005). For example, *Combretum zeyheri* was preferred on a local scale during the dry season but became less important when the resource availability scale enlarged to the landscape level. The shift of forage preference as spatial scale increases is presumably associated with aggregates of habitat units within a heterogeneous landscape, which in turn affects estimates of resource availability (Li and Reynolds 1993; Bissonetie et al. 1997), and consequently, resource selection decisions. Nevertheless, giraffes at the landscape scale exhibited similar foraging selection patterns, with only two more forage species preferred at least once across the season, compared to the local scale. The detected slight differences in forage selection patterns may be attributed to giraffes' relatively limited seasonal migrations in savanna landscapes (Pellew 1984b), making only a portion of forage options accessible to giraffes. The multispatial scaled analyses for forage-procuring strategies that were used in this study provide a reliable means of comparing use versus available estimates at local and landscape scales of selection (Bissonetie et al. 1997; Mysterud et al. 1999; Anderson et al. 2005; Fortin et al. 2005; Boyce 2006; Kotliar and Wiens 2013), which eventually leads to judicious management decisions (Andren 1994; Johnson et al. 2002).

Our observed high and year-round preference for the native shrub *D. cinerea* at both local and landscape scales might be linked to *D. cinerea*'s high nutritional quality (Pellew 1984b; Mlambo et al. 2004; Smith et al. 2005; Dagg 2014; Tjelele et al. 2014) despite the belief that this plant is unpalatable. Further, *D. cinerea*'s gradual loss of leaves followed by quick recovery upon the onset of the wet season (Sauer 1983) might ensure that giraffes receive a constant supply of this food resource throughout the year. Hence, despite the negative attitudes of livestock managers toward *D. cinerea* in a rangeland ecosystem, this plant species might be contributing significantly to the quality and quantity of food for the full suite of savanna browsers (Reyes et al. 2010; Liu et al. 2015).

Conservation implications.—Giraffes in the Tarangire Ecosystem selected a broad array of woody plant species from which some were preferred, while others were avoided. Forage selection in giraffes was influenced primarily by spatial and temporal changes in the quantities and, presumably, qualities of forage species at both local and landscape scales. Giraffes also

showed a strong preference for the native shrub *D. cinerea* at both local and landscape scales. *Dichrostachys cinerea* is a fast-growing nutritive shrub, well adapted in the study area, palatable to giraffes, and resistant to strong browsing pressure, providing giraffe with a constant supply of this food resource throughout the year. The removal of this species for the purpose of maintaining grazing lawns for livestock might negatively impact browsing wildlife. Therefore, management that is focused on benefiting grazing livestock by removal of encroaching woody plant species may have unintended consequences to browsing species such as giraffes that feed extensively on these food resources.

ACKNOWLEDGMENTS

The German Academic Exchange Service (DAAD), Rufford Foundation (Small Grants for Nature), Save The Giraffes, Cleveland Zoological Society (CZS), Wild Nature Institute, American Society of Zoologists, Explorers Club Grants, and IDEA WILD are sincerely acknowledged for their financial support. We thank Mr. Stoni, Erasto, Oscar, Monsoni, Alfred, and other field assistants for helping with data collection and plant identification. We are grateful to the Tanzania Commission for Science and Technology (COSTECH), Tanzania Wildlife Research Institute (TAWIRI), Tanzania Wildlife Authority (TAWA), Randilen Wildlife Management Area, Manyara Ranch Conservancy, African Wildlife Foundation (AWF), and Tanzania National Parks (TANAPA) for permitting this study.

SUPPLEMENTARY DATA

Supplementary data are available at *Journal of Mammalogy* online.

Supplementary Data SD1.—Woody plant species consumed by Masai giraffes (*Giraffa camelopardalis tippelskirchi*) in the Tarangire Ecosystem of Tanzania in 2019, and their relative proportions in the diet (P_j) over the entire year and split up into the dry and wet season based on direct foraging observations ($n = 3,728$) during scan sampling.

Supplementary Data SD2.—Woody plant species encountered in vegetation sampling in 2019 in Tarangire Ecosystem, Tanzania.

Supplementary Data SD3.—Masai giraffe (*Giraffa camelopardalis tippelskirchi*) local-scale forage selection ratios (\hat{W}), and their Bonferroni confidence intervals (CIs \hat{W}), and P -values for seasonal selection differences for 38 woody plant species in the Tarangire Ecosystem, 2019.

Supplementary Data SD4.—Masai giraffe (*Giraffa camelopardalis tippelskirchi*) landscape-scale forage selection ratios (\hat{W}), their Bonferroni confidence intervals (CIs \hat{W}), and seasonal selection differences for 38 woody plant species in the Tarangire Ecosystem, 2019.

Supplementary Data SD5.—Masai giraffe feeding on *Dichrostachys cinerea* in Tarangire National Park in 2019.

LITERATURE CITED

Anderson D.P., Turner M.G., Forester J.D., Zhu J., Boyce M.S., Beyer H., Stowell L. 2005. Scale-dependent summer resource

- selection by reintroduced elk in Wisconsin, USA. *Journal of Wildlife Management* 69:298–310.
- Andren H. 1994. Effects of habitat fragmentation on birds and mammals of suitable habitat. *Oikos* 71:355–366.
- Arthur S.M., Manly B.F.J., McDonald L.L., Garner G.W. 1996. Assessing habitat selection when availability changes. *Ecology* 77:215–227.
- Banda T., Mwangulango N., Meyer B., Schwartz M.W., Mbago F., Sungula M., Caro T. 2008. The woodland vegetation of the Katavi-Rukwa ecosystem in western Tanzania. *Forest Ecology and Management* 255:3382–3395.
- Belayneh A., Tessema Z.K. 2017. Mechanisms of bush encroachment and its inter-connection with rangeland degradation in semi-arid African ecosystems. *Journal of Arid Lands* 9:299–312.
- Bercovitch F.B., Berry P.S.M. 2013. Herd composition, kinship and fission–fusion social dynamics among wild giraffe. *African Journal of Ecology* 15:206–216.
- Berry P. S. M., Bercovitch F. B. 2016. Seasonal and geographical influences on the feeding ecology of giraffes in the Luangwa Valley, Zambia. *African Journal of Ecology* 55:80–90.
- Beyer E. Jr., Haufler J.B. 1994. Diurnal versus 24-hour sampling of habitat use. *Journal of Wildlife Management* 58:178–180.
- Birhane E., Treydte A.C., Eshete A., Solomon N., Hailemariam M. 2017. Can rangelands gain from bush encroachment? Carbon stocks of communal grazing lands invaded by *Prosopis Juliflora*. *Journal of Arid Environments* 141:60–67.
- Bissonette J.A., Harrison D.J., Hargis C.D., Chapin T.G. 1997. The influence of spatial scale and scale-sensitive properties on habitat selection by American marten. In: Bissonette J.A., editor. *Wildlife and landscape ecology*. Springer, New York, USA; p. 368–385.
- Bolger D.T., Newmark W.D., Morrison T.A., Doak D.F. 2008. The need for integrative approaches to understand and conserve migratory ungulates. *Ecology Letters* 11:63–77.
- Bolger D., Ogutu J., Strauss M., Lee D., Muneza A., Fennessy J., Brown D. 2019. *Giraffa camelopardalis* ssp. *tippelskirchi*, Masai giraffe. The IUCN Red List of threatened species. 2019 version 3.1. www.iucnredlist.org. Accessed 23 September 2019.
- Bond M.L., Lee D.E., Ozgul A., König B. 2019. Fission–fusion dynamics of a megaherbivore are driven by ecological, anthropogenic, temporal, and social factors. *Oecologia* 191:335–347.
- Boyce M.S. 2006. Scale for resource selection functions. *Diversity and Distributions* 12:269–276.
- Bryson-Morrison N., Tzanopoulos J., Matsuzawa T., Humle T. 2017. Activity and habitat use of chimpanzees (*Pan troglodytes verus*) in the anthropogenic landscape of Bossou, Guinea, West Africa. *International Journal of Primatology* 38:282–302.
- Caister L.E., Shields W.M., Gosser A. 2003. Female tannin avoidance: a possible explanation for habitat and dietary segregation of giraffes (*Giraffa camelopardalis peralta*) in Niger. *African Journal of Ecology* 41:201–210.
- Chytry M., Tichy L., Hennekens S.M., Schaminee J.H.J. 2013. Assessing vegetation change using vegetation-plot databases: a risky business. *Applied Vegetation Science* 17:32–41.
- Dagg A.I. 2014. Giraffe. Biology, behaviour, and conservation. Vol. 53. Cambridge University Press, New York, USA.
- Dalle G., Maass B.L., Isselstein J. 2006. Encroachment of woody plants and its impact on pastoral livestock production in the Borana lowlands, southern Oromia, Ethiopia. *African Journal of Ecology* 44:237–246.
- Devine A.P., McDonald R.A., Quaipe T., Maclean I.M.D. 2017. Determinants of woody encroachment and cover in African savannas. *Oecologia* 183:939–951.

- Dumont B., Rook A.J., Coran C., Rover K.U. 2007. Effects of live-stock breed and grazing intensity on biodiversity and production in grazing systems. *Grass and Forage Science* 62:159–171.
- Fortin D., Morales J.M., Boyce M.S. 2005. Elk winter foraging at fine-scale in Yellowstone National Park. *Oecologia* 145:335–343.
- Galanti V., Preatoni D., Martinoli A., Wauters L.A., Tosi G. 2006. Space and habitat use of the African elephant in the Tarangire–Manyara Ecosystem, Tanzania: implications for conservation. *Mammalian Biology* 71:99–14.
- Grady J.M., Hoffmann W.A. 2012. Caught in a fire trap: recurring fire creates stable size equilibria in woody resprouters. *Ecology* 93:2052–2060.
- Hudak A.T., Wessman C.A. 1998. Textural analysis of historical aerial photography to characterize woody plant encroachment in South African savanna. *Remote Sensing of Environment* 66:317–330.
- Johnson D.H. 1980. The comparison of usage and availability measurements for evaluating resource preference. *Ecology* 61:65–71.
- Johnson C.J., Parker K.L., Heard D.C., Gillingham M.P. 2002. A multiscale behavioral approach to understanding the movements of woodland caribou. *Ecological Applications* 12:1840–1860.
- Kahurananga J., Silkiluwasha F. 1997. The migration of zebra and wildebeest between Tarangire National Park and Simanjoro plains, Northern Tanzania, in 1972 and recent trends. *African Journal of Ecology* 35:179–185.
- Kotliar N.B., Wiens J.A. 2013. Multiple scales of patchiness and patch structure: a hierarchical for the study of heterogeneity framework. *Oikos* 59:253–260.
- Lee D.E., Bolger D.T. 2017. Movements and source-sink dynamics of a Masai giraffe metapopulation. *Population Ecology* 59:157–168.
- Lee D.E., Bond M.L. 2016. Precision, accuracy, and costs of survey methods for giraffe *Giraffa camelopardalis*. *Journal of Mammalogy* 97:940–948.
- Lee D.E., Bond M.L., Kissui B.D.M., Kiwango Y.A., Bolger D.T. 2016. Spatial variation in giraffe demography: a test of 2 paradigms. *Journal of Mammalogy* 97:1015–1025.
- Li H., Reynolds J.F. 1993. A new contagion index to quantify spatial patterns of landscapes. *Landscape Ecology* 8:155–162.
- Lindgren P.M.F., Sullivan T.P. 2001. Influence of alternative vegetation management treatments on conifer plantation attributes: abundance, species diversity, and structural diversity. *Forest Ecology and Management* 142:163–182.
- Liu J., Feng C., Wang D., Wang L., Wilsey B.J., Zhong Z. 2015. Impacts of grazing by different large herbivores in grassland depend on plant species diversity. *Journal of Applied Ecology* 52:1053–1062.
- Ludwig F. 2001. Tree-grass interactions on an east African savanna: the effects of competition, facilitation, and hydraulic lift. *Tropical Resource Management Papers* 39:144.
- Ludwig F., De Kroon H., Prins H.H.T. 2008. Impacts of savanna trees on forage quality for a large African herbivore. *Oecologia* 155:487–496.
- Mahenya O., Mathisen K.M., Andreassen H.P., Skarpe C. 2016. Hierarchical foraging by giraffe in a heterogeneous savannah, Tanzania. *African Journal of Ecology* 54:136–145.
- Manly B.F.J., McDonald L.L., Thomas D.L., McDonald T.L., Erickson W.P. 2002. Resource selection by animals: statistical design and analysis for field studies. 2nd ed. Kluwer Academic Publishers, New York, USA.
- Martin P., Bateson P. 1993. Measuring behaviour: an introductory guide. 2nd ed. Cambridge University Press, Cambridge, United Kingdom.
- Maslin B.R. 2008. Generic and subgeneric names in acacia following retypification of the genus. *Mulleria* 26:7–9.
- Mlambo V., Smith T., Owen E., Mould F.L., Sikosana J.L.N., Mueller-Harvey I. 2004. Tanniniferous *Dichrostachys cinerea* fruits do not require detoxification for goat nutrition: in sacco and in vivo evaluations. *Livestock Production Science* 90:135–144.
- Msoffe F.U., Kifugo S.C., Said M.Y., Neselle M.O., van Gardingen P., Reid R.S., Ogutu J.O., Herero M., de Leeuw J. 2011. Drivers and impacts of land-use change in the Maasai steppe of Northern Tanzania: an ecological, social, and political analysis. *Journal of Land Use Science* 6:261–281.
- Mysterud A., Lian L., Hjernmann D. 1999. Scale-dependent trade-offs in foraging by European roe deer (*Capreolus capreolus*) during winter. *Canadian Journal of Zoology* 77:1486–1493.
- Neu C.W., Byers C.R., Peek J.M. 1974. A technique for analysis of utilization-availability data. *Journal of Wildlife Management* 38:541–545.
- Newmark W.D., Manyaza D.N., Gamassa D.M., Sariko H.I. 1994. The conflict between wildlife and local people living adjacent to protected areas in Tanzania: human density as a predictor. *Conservation Biology* 8:249–255.
- Njagi D. 2019. In east Africa, spread of sickle bush drives conflicts with wildlife, March 2019. <https://news.mongabay.com/2019/03/in-east-africa-spread-of-sickle-bush-drives-conflict-with-wildlife/>. Accessed 15 July 2019.
- Orians G.H., Wittenberge J.F. 1991. Spatial and temporal scales in habitat selection. *American Naturalist* 137:29–49.
- Parker D.M., Bernard R.T.F. 2005. The diet and ecological role of giraffe (*Giraffa camelopardalis*) introduced to the Eastern Cape, South Africa. *Journal of Zoology* 267:203–210.
- Parke D.M., Bernard R.T.F., Colvin S.A. 2003. The diet of a small group of extralimital giraffe. *African Journal of Ecology* 41:245–253.
- Pellew R.A. 1983. The giraffe and its food resource in the Serengeti: response of the giraffe population to changes in the food supply. *African Journal of Ecology* 21:241–267.
- Pellew R.A. 1984a. Food consumption and energy budgets of the giraffe. *Journal of Applied Ecology* 21:141–159.
- Pellew R.A. 1984b. The feeding ecology of a selective browser, the giraffe (*Giraffa camelopardalis tippelskirchi*). *Journal of Zoology* 202:57–81.
- Peterson D.D. 1978. Seasonal distributions and interactions of cattle and wild ungulates in Masailand, Tanzania. M.S. thesis, Virginia Polytechnic Institute, and State University, Blacksburg, Virginia.
- Pratt D.J., Greenway P.J., Gwynne M.D. 1966. A classification of East African rangeland, with an appendix on terminology. *Journal of Applied Ecology* 3:369–382.
- Proll J., Petzke K.J., Ezeagu I.E., Metges C.C. 2018. Nutrient requirements and interactions low nutritional quality of unconventional tropical crop seeds in rats. *Journal of Nutrition* 128:2014–2022.
- Reyes G.P., Kneeshaw D., Grandpre L.D., Leduc A. 2010. Changes in woody vegetation abundance and diversity after natural disturbances causing different levels of mortality. *Journal of Vegetation Science* 21:406–417.
- Roques K.G., O'Connor T.G., Watkinson A.R. 2001. Dynamics of shrub encroachment in an African savanna: relative influences of fire, herbivory, rainfall, and density dependence. *Journal of Applied Ecology* 38:268–280.
- Sauer J.J.C. 1983. Food selection by giraffes in relation to changes in chemical composition of the leaves. *South African Journal of Animal Science* 3:1980–1983.
- Sauer J.J.C., Theron G.K., Skinner J.D. 1977. Food preferences of giraffe, *Giraffa camelopardalis* in the arid bushveld of the Western Transvaal. *South African Journal of Wildlife Research* 7:53–59.

- Shiple L.A. 2007. The influence of bite size on foraging at larger spatial and temporal scales by mammalian herbivores. *Oikos* 116:1964–1974.
- Shorrocks B., Bates W. 2015. *The biology of African savannahs*. 2nd ed. Oxford University Press, Oxford, United Kingdom.
- Sikes R.S.; The Animal Care and Use Committee of the American Society of Mammalogists. 2016. Guidelines of the American Society of Mammalogists for the use of wild mammals in research and education. *Journal of Mammalogy* 97:663–688.
- Smit I.P.J., Prins H.H.T. 2015. Predicting the effects of woody encroachment on mammal communities, grazing biomass, and fire frequency in African savannas. *PLoS ONE* 10:e0137857.
- Smith T., Mlambo V., Sikosana J.L.N., Maphosa V., Mueller-Harvey I., Owen E. 2005. *Dichrostachys cinerea* and *Acacia nilotica* fruits as dry season feed supplements for goats in a semi-arid environment: summary of a DFID funded project in Zimbabwe. *Animal Feed Science and Technology* 122:149–157.
- Stoner C., Caro T., Mduma S., Mlingwa C., Sabuni G., Borner M. 2007. Assessment of effectiveness of protection strategies in Tanzania based on a decade of survey data for large herbivores. *Conservation Biology* 21:635–646.
- Tjelele J., Ward D., Dziba L. 2014. Diet quality modifies germination of *Dichrostachys cinerea* and *Acacia nilotica* seeds fed to ruminants. *Rangeland Ecology and Management* 67:423–428.
- Van de Vijver C.A.D.M., Foley C.A., Olff H. 1999. Changes in the woody component of an east African savanna during 25 years. *Journal of Tropical Ecology* 15:545–564.
- Whittingham M.J., Swetnam R.D., Wilson J.D., Chamberlain D.E., Freckleton R.P. 2005. Habitat selection by yellowhammers *Emberiza citrinella* on lowland farmland at two spatial scales: implications for conservation management. *Journal of Applied Ecology* 42:270–280.
- Wiens J.A. 1981. Scale problems in avian censusing. *Studies in Avian Biology* 6513:513–521.
- Woodward A., Coppock D.L. 1995. Role of plant defense in the utilization of native browse in Southern Ethiopia. *Agroforestry Systems* 32:147–161.
- Zarovali M.P., Yiakoulaki M.D., Papanastasis V.P. 2007. Effects of shrub encroachment on herbage production and nutritive value in semi-arid Mediterranean grasslands. *Grass and Forage Science* 62:355–363.

Submitted 7 June 2020. Accepted 18 January 2022.

Associate Editor was Kevin Monteith.